

Exterior Differential Systems for Yang–Mills Theories^{*}

Frank B. ESTABROOK

Jet Propulsion Laboratory, California Institute of Technology, Pasadena, CA 91109, USA

E-mail: frank.b.estabrook@jpl.nasa.gov

Received July 18, 2008, in final form September 03, 2008; Published online September 12, 2008

Original article is available at <http://www.emis.de/journals/SIGMA/2008/063/>

Abstract. Exterior differential systems are given, and their Cartan characters calculated, for Maxwell and $SU(2)$ -Yang–Mills equations in dimensions from three to six.

Key words: exterior differential systems; Cartan characters; Maxwell equations; $SU(2)$ -Yang–Mills equations

2000 Mathematics Subject Classification: 58A15

In memory of my friend and colleague Hugo D. Wahlquist [1930–2008]

1 Introduction

In this work Exterior Differential Systems (EDS) are formulated for abelian and $SU(2)$ -gauge theories in 3 to 6 dimensions, beginning with Maxwell vacuum electrodynamics and $SU(2)$ -Yang–Mills theory in 4 dimensions. The generalization to larger groups and n dimensions becomes apparent. The results are new in that the Cartan integer character tables are explicitly calculated, showing the EDS' to be well posed field theories. Cartan characters yield the numbers of evolution equations, of first order (second order, etc.), of constraint equations (or integrability conditions), and of gauge degrees of freedom; well posed EDS' allow systematic derivation of isovectors and similarity solutions, conservation laws and prolongations.

An EDS for vacuum Maxwell theory in 4 dimensions has been set on a 14 dimensional manifold [1, 2, 3] with variables A_i , $F_{ij} = -F_{ji}$, x^i , $i = 1, 2, 3, 4$. Its generalization to Yang–Mills $SU(2)$, in terms of 34 variables A_i^a , $F_{ij}^a = -F_{ji}^a$, x^i , $i = 1, 2, 3, 4$, $a = 1, 2, 3$ was also partially formulated in the first two of these references. An analogous EDS for vacuum general relativity was more recently found, and Cartan characters and conservation laws calculated for both it and vacuum Maxwell theory [4, 3]. These EDS' are generated by sets of “torsion” 2-forms and dynamic 3-forms, and these factor the terms of a Cartan–Poincaré (or multisymplectic) 5-form $d\Lambda$ expressing the underlying variational principle. The generalized EDS' in n dimensions to be given in Sections 2 and 3 will follow this structure, using torsion 2-forms and $(n-1)$ -forms.

It should be noted that this approach to finding a Cartan n -form Λ , or Lagrangian density, for a field theory differs from one that constructs a Cartan–Poincaré form $d\Lambda$ using “contact” 1-forms and dynamic n -forms [5, 6]. That approach has been carefully worked out as “multi-contact” geometry [7]. In 4-dimensional Maxwell theory, e.g., the multicontact EDS is set on a 24 dimensional manifold, spanned by four potential fields A^i , 16 fields A_j^i and the x^i . The Cartan characteristic integer table is quite different.

The Cartan forms and EDS' in this work were manipulated and checked algebraically using the MATHEMATICA suite of programs AVF (Algebra Valued Forms) written by the late

^{*}This paper is a contribution to the Special Issue “Élie Cartan and Differential Geometry”. The full collection is available at <http://www.emis.de/journals/SIGMA/Cartan.html>

H.D. Wahlquist [8]. The integer characters were calculated using his Monte Carlo program `characters.m`, which assigns random integer values to unknown components of successively calculated vectors spanning integral manifolds of an EDS, vectors satisfying the nested linear equations of Cartan–Kuranishi integration theory. After three or so of these are calculated, using integer arithmetic, the new vector components solved for become rational numbers with scores of digits, and the consequent calculation of the ranks of the resulting matrices (and so the integer characters) is absolutely robust. (In principle of course this program could be rewritten to use the first few hundred primes instead of random integers, making the calculation of characters exact.)

In 2003 Wahlquist circulated on CD a final version of his principal MATHEMATICA notebooks, with documentation and demonstrations, as “AVF Programs”. This is now available from the SIGMA web-site: <http://www.emis.de/journals/SIGMA/2008/063/Wahlquist2003.zip>.

2 Maxwell and Yang–Mills

We will be generalizing the EDS and Cartan form for the vacuum Maxwell equations; these are most conveniently written using a 2-form $F = \frac{1}{2}F_{ij}dx^i \wedge dx^j$ and a dual 2-form $*F = \frac{1}{4}F_{ij}dx^k \wedge dx^l \epsilon^{ij}_{..kl}$ involving the fields and a 1-form $A = A_i dx^i$ coding the potentials. $i, j = 1, \dots, 4$, and indices are raised/lowered with the Lorentz metric. The EDS is generated [3] by a 2-form $\theta = dA - F$, its closure $d\theta$ and another 3-form $\psi = -d*F$. The Cartan characters are $s_0 = 0$, $s_1 = 1$, $s_2 = 3$, $s_3 = 5$, $s_4 = 1$. The genus $g = 4$ (the dimension of the maximal regular integral manifold, or solution), and the dx^i are in involution (so independent variables in the PDE.) We summarize this well-posed structure by writing $14[0,1,3,5]4+1$. The last integer, s_4 , is the number of arbitrary functions or gauge freedom in a solution. There are no Cauchy vectors, so Cartan’s test for well-posedness requires that the first integer equals the sum of the rest.

The Cartan–Poincaré form for this EDS is

$$d\Lambda = \theta \wedge \psi. \quad (1)$$

That this outer product is indeed closed as we have written follows from a direct computation that $d(\theta \wedge \psi) = 0$. This in turn is true because

$$dF \wedge d*F = 0$$

which is nothing else than $dF_{ij} \wedge dF_{ij} = 0$. Integrating equation (1) by parts, we take

$$\Lambda = -*F \wedge dA + \frac{1}{2}F \wedge *F.$$

This Cartan form corresponds to the Lagrangian density for Maxwell theory used by field theorists [9, page 98, equation (9.13)].

The generalization to Yang–Mills theory in 4 dimensions is of course well known. Its form is nicely exposed when written as an EDS. Structure constants of an m -dimensional Lie group γ_{bc}^a , $a, b, \dots = 1, \dots, m$ are used, taken in a basis where these are completely antisymmetric ($\gamma_{bc}^a = -\gamma_{cb}^a = \gamma_{ca}^b = \gamma_{ab}^c$) and we take $\gamma_{23}^1 = 1$. Quadratically nonlinear terms are introduced consistently coupling indexed variables A_i^a and F_{ij}^a . In the following we consider only the case $m = 3$, which is $SU(2)$. We define 1-forms and 2-forms

$$\begin{aligned} A^a &= A_i^a dx^i, \\ F^a &= \frac{1}{2}F_{ij}^a dx^i \wedge dx^j, \\ *F_a &= \frac{1}{4}F_{ij}^a dx^k \wedge dx^l \epsilon^{ij}_{..kl} \end{aligned}$$

and now set

$$\begin{aligned}\theta^a &= dA^a - F^a + \frac{1}{8}\gamma_{bc}^a A^b \wedge A^c, \\ \psi_a &= -d * F_a - \frac{1}{4}\gamma_{ab}^c A^b \wedge *F_c.\end{aligned}$$

The EDS in 34 variables is generated by the three 2-forms θ^a , three 3-forms $d\theta^a$ and three additional 3-forms ψ_a . That this EDS is closed follows from calculating

$$d\psi_a = -\frac{1}{4}\gamma_{ab}^c A^b \wedge \psi_c - \frac{1}{4}\gamma_{ab}^c \theta^b \wedge *F_c,$$

so $d\psi_a$ are in the EDS (the essential identity in this is $\gamma_{ab}^c *F_c \wedge F^b = 0$).

The Cartan–Poincaré form is now

$$d\Lambda = \theta^a \wedge \psi_a$$

and it can be explicitly integrated to take

$$\Lambda = - *F_a \wedge dA^a + \frac{1}{2}F^a \wedge *F_a - \frac{1}{8}\gamma_{ab}^c A^a \wedge A^b \wedge *F_c.$$

A possibly new result is a calculation of the Cartan character table. We find

$$34[0, 3, 9, 15]4 + 3$$

showing this theory to be well-posed, with 3 degrees of gauge freedom.

3 Generalization to other dimensions

All the equations of Section 2 can be taken as they stand in any number of spacetime dimensions, $i, j = 1, \dots, n$. We need only redefine the dual field form as an $(n - 2)$ -form

$$*F_a = \frac{1}{2(n-2)!} F_{ij}^a dx^k \wedge dx^l \wedge \dots \wedge dx^n \epsilon_{..kl\dots n}^{ij}.$$

The ψ_a are now $(n - 1)$ -forms. We have explicitly verified this generalization, and calculated the Cartan characters for theories through $n = 6$, showing the EDS' to be well-posed. These results are summarized in Table 1, and the pattern for higher dimensions is apparent.

Table 1. Calculated Cartan characters for Maxwell and $SU(2)$ -Yang–Mills theories in n dimensions. The characters $s_0, \dots, s_{n-1}$ are given in brackets [], followed by n . The preceding number is the dimension of the manifold where the EDS is set. The last number is s_n , the gauge freedom.

Maxwell characters in n dimensions

$$\begin{aligned}9[0, 2, 3]3 + 1 \\ 14[0, 1, 3, 5]4 + 1 \\ 20[0, 1, 2, 4, 7]5 + 1 \\ 27[0, 1, 2, 3, 5, 9]6 + 1\end{aligned}$$

$SU(2)$ -Yang–Mills characters in n dimensions

$$\begin{aligned}21[0, 6, 9]3 + 3 \\ 34[0, 3, 9, 15]4 + 3 \\ 50[0, 3, 6, 12, 21]5 + 3 \\ 69[0, 3, 6, 9, 15, 27]6 + 3\end{aligned}$$

Acknowledgements

This work was performed while I held a visiting appointment at the Jet Propulsion Laboratory, California Institute of Technology.

References

- [1] Estabrook F.B., Hermann R., Wahlquist H.D., Appendix D, in Hermann R., Yang–Mills, Kaluza–Klein and the Einstein Program, *Interdisciplinary Mathematics*, Vol. XIX, Math Sci Press, Brookline, Mass., 1978.
- [2] Hermann R., Differential form methods in the theory of variational systems and Lagrangian field theories, *Acta Appl. Math.* **12** (1998), 35–78.
- [3] Estabrook F.B., Conservation laws for vacuum tetrad gravity, *Classical Quantum Gravity* **23** (2006), 2841–2848, gr-qc/0508081.
- [4] Estabrook F.B., Mathematical structure of tetrad equations for vacuum relativity, *Phys. Rev. D* **71** (2005), 044004, 5 pages, gr-qc/0411029.
- [5] Hojman S., Problem of the identical vanishing of Euler–Lagrange derivatives in field theory, *Phys. Rev.* **27** (1983), 451–453.
- [6] Betounes D.E., Extension of the classical Cartan form, *Phys. Rev. D* **29** (1984), 599–606.
- [7] Bryant R., Griffiths P., Grossman D., Exterior differential systems and Euler–Lagrange partial differential equations, The University of Chicago Press, Chicago – London, 2003.
- [8] Wahlquist H.D., Monte Carlo calculation of Cartan characters, in Proceedings of the International Conference of Aspects of General Relativity and Mathematical Physics, Editors N. Breton, R. Capovilla and T. Matos, Ediciones CINVESTAV, Mexico DF, 1994, 168–174.
- [9] Schwinger J., DeRaad L.L. Jr., Milton K.A., Tsai W.-Y., Classical electrodynamics, Westview Press, Boulder CO, 1998.